

TOM J. POUSTI, MD, F.A.C.S.
PLASTIC AND RECONSTRUCTION SURGERY

**INFORMED CONSENT FOR LIPOSUCTION
(SUCTION-ASSISTED LIPECTOMY SURGERY)
(ULTRASOUND-ASSISTED LIPECTOMY SURGERY)**

Instructions

This is an informed-consent document that has been prepared to help inform you concerning suction-assisted lipectomy and ultrasound-assisted lipectomy (“liposuction”) surgery, its risks, and alternative treatment.

It is important that you read this information carefully and completely. Please initial each page, indicating that you have read the page and sign the consent for surgery as proposed by your plastic surgeon.

Introduction

Liposuction is a surgical technique to remove unwanted deposits of fat from specific areas of the body, including the face and neck, upper arms, trunk, abdomen, buttocks, hips and thighs, and the knees, calves, and ankles. This is not a substitute for weight reduction, but a method for removing localized deposits of fatty tissue that do not respond to diet or exercise. Liposuction may be performed as a primary procedure for body contouring or may be combined with other surgical techniques such as facelift, abdominoplasty, or thigh lift procedures to tighten loose skin and supporting structures.

The best candidates for liposuction are individuals of relatively normal weight who have excess fat in particular body areas. Having firm, elastic skin will result in a better final contour after liposuction. Hanging skin will not reshape itself to the new contours and may require additional surgical techniques to remove and tighten excess skin. Body contour irregularities due to structures other than fat cannot be improved by this technique. Liposuction itself will not improve areas of dimpled skin known as “cellulite”.

Suction-assisted lipectomy surgery is performed by using a hollow metal surgical instrument known as a cannula that is inserted through small skin incision(s) and is passed back and forth through the area of fatty deposit. The cannula is attached to a vacuum source, which provides the suction needed to remove the fatty tissue.

In some situations, a special cannula may be used that emits ultrasonic energy to break down fatty deposits. This technique is known as **ultrasonic-assisted lipectomy**. Depending on your needs, your surgeon may recommend suction-assisted lipectomy alone, or in combination with ultrasound-assisted lipectomy.

There are a variety of different techniques used by plastic surgeons for liposuction and care following surgery. Liposuction may be performed under local or general anesthesia. **Tumescent liposuction technique** involves the infiltration of fluid containing dilute local anesthetic and epinephrine into areas of fatty deposits. This technique can reduce discomfort at the time of surgery, as well as reduce post-operative bruising.

Support garments and dressings are worn to control swelling and promote healing. Your surgeon may recommend that you make arrangements to donate a unit of your own blood that would be used if a blood transfusion were necessary after surgery.

Alternative Treatment

Alternative forms of management consist of not treating the areas of fatty deposits. Diet and exercise regimens may be of benefit in the overall reduction of excess body fat.

Direct removal of excess skin and fatty tissue may be necessary in addition to liposuction in some patients. Risks and potential complications are associated with alternative forms of treatment that involve surgery.

Risks of Liposuction Surgery

Every surgical procedure involves a certain amount of risk, and it is important that you understand the risks involved with liposuction. An individual’s choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. Although the majority of patients do not experience these complications, you

should discuss each of them with your plastic surgeon to make sure you understand the risks, potential complications, and consequences of liposuction.

Patient Selection

Individuals with poor skin tone, medical problems, obesity, or unrealistic expectations may not be candidates for liposuction.

Bleeding

It is possible, though unusual, to have a bleeding episode during or after surgery. Should post-operative bleeding occur, it may require emergency treatment to drain accumulated blood or require a blood transfusion. Do not take any aspirin or anti-inflammatory medications for seven days before surgery, as this may increase the risk of bleeding.

Infection

An infection is quite unusual after this type of surgery. Should an infection occur, treatment including antibiotics or additional surgery may be necessary.

Skin Scarring

Although, good wound healing after a surgical procedure is expected, abnormal scars may occur within the skin and deeper tissues. In rare cases, abnormal scars may result. Scars may be unattractive and of a different color than surrounding skin. Additional treatments, including surgery, may be needed to treat abnormal scarring.

Change in Skin Sensation

A temporary decrease in skin sensation after liposuction may occur. This usually resolves over a period of time. Diminished (or complete loss of skin sensation) infrequently occurs and may not totally resolve.

Skin Discoloration/Swelling

Skin discoloration and swelling normally occurs following liposuction. In rare situations, swelling and skin discoloration may persist for long periods of time. Permanent skin discoloration is rare.

Skin Contour Irregularities

Contour irregularities and depressions in the skin may occur after liposuction. Visible and palpable wrinkling of skin can occur. Additional treatments including surgery may be necessary to treat skin contour irregularities following Liposuction.

Asymmetry

It may not be possible to achieve symmetrical body appearance from liposuction surgery. Factors such as skin tone, bony prominence, and muscle tone may contribute to normal asymmetry in body features.

Seroma

Fluid accumulations infrequently occur in areas where liposuction has been performed. Additional treatments or surgery to drain accumulations of fluid may be necessary.

Long Term Effects

Subsequent alterations in body contour may occur as the result of aging, weight loss or gain, pregnancy, or other circumstances not related to liposuction.

Tumescent Liposuction

There is the possibility that large volumes of fluid containing dilute local anesthetic drugs and epinephrine that is injected into fatty deposits during surgery may contribute to fluid overload or systemic reaction to these medications. Additional treatment including hospitalization may be necessary.

Other

You may be disappointed with the results of surgery. Infrequently, it is necessary to perform additional surgery to improve your results.

Surgical Anesthesia

Both local and general anesthesia involve risk. There is the possibility of complication, injury, and even death from all forms of surgical anesthesia or sedation.

Allergic Reaction

In rare cases, local allergies to tape, suture material, or topical preparations have been reported. Systemic reactions, which are more serious, may occur due to drugs used during surgery and prescription medicines. Allergic reactions may require additional treatment.

Surgical Shock

In rare circumstances, liposuction can cause severe trauma, particularly when multiple or extensive areas are suctioned at one time. Although serious complications are infrequent, infections or excessive fluid loss can lead to severe illness and even death. Should surgical shock occur after liposuction, hospitalization and additional treatment would be necessary.

Pulmonary Complications

Fat embolism syndrome occurs when fat droplets are trapped in the lungs. This is a very rare and possible fatal complication of suction-assisted lipectomy. Should fat embolism or other pulmonary complications occur, additional treatment including hospitalization may be necessary.

Skin Loss

Skin loss is very rare after liposuction. Additional treatments including surgery may be necessary.

Chronic Pain

Chronic pain and discomfort following liposuction is very rare.

Ultrasound-Assisted Lipectomy

Risks associated with the use of this technique include the above mentioned risks and the following specific risks:

BURNS

Ultrasonic energy may produce burns and tissue damage either at the location where the cannula is inserted into the skin or in other areas if the cannula touches the undersurface of the skin for prolonged periods of time. If burns occur, additional treatment and surgery may be necessary.

CANNULA FRAGMENTATION

Ultrasonic energy produced within the cannula may cause disintegration (fragmentation) of the surgical instrument. The occurrence and effect of this is unpredictable. Should this occur, additional treatment, including surgery, may be necessary.

UNKNOWN RISKS

The long-term effect on tissue and organs to exposure too short-duration, high-intensity ultrasonic energy is unknown. There is the possibility that additional risk factors of ultrasound-assisted lipectomy may be discovered.

Health Insurance

Most health insurance companies exclude coverage for cosmetic surgical operations such as liposuction or any complications that might occur from surgery. Please carefully review your health insurance subscriber-information pamphlet.

Additional Surgery Necessary

There are many variable conditions in addition to risk and potential surgical complications that may influence the long term result from liposuction. Even though risks and complications occur infrequently, the risks cited are particularly associated with liposuction. Other complications and risks can occur but are even more uncommon. Should complications occur, additional surgery or other treatments may be necessary. The practice

of medicine and surgery is not an exact science. Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained.

Financial Responsibilities

The cost of surgery involves several charges for the services provided. The total includes fees charged by your doctor, the cost of surgical supplies, anesthesia, laboratory tests, and possible outpatient hospital charges, depending on where the surgery is performed. Depending on whether the cost of surgery is covered by an insurance plan, you will be responsible for necessary co-payments, deductibles, and charges not covered. Additional costs may occur should complications develop from the surgery. Secondary surgery or hospital day-surgery charges involved with revisionary surgery would also be your responsibility.

Disclaimer

Informed-consent documents are used to communicate information about the proposed surgical treatment of a disease or condition along with disclosure of risks and alternative forms of treatment(s). The informed-consent process attempts to define principles of risk disclosure that should generally meet the needs of most patients in most circumstances.

However, informed consent documents should not be considered all inclusive in defining other methods of care and risks encountered. Your plastic surgeon may provide you with additional or different information, which is based on all the facts in your particular case and the state of medical knowledge.

Informed-consent documents are not intended to define or serve as the standard of medical care. Standards of medical care are determined on the basis of all of the facts involved in an individual case and are subject to change as scientific knowledge and technology advance and as practice patterns evolve.

It is important that you read the above information carefully and have all of your questions answered before signing the consent on the next page.

CONSENT FOR SURGERY/PROCEDURE or TREATMENT

I hereby authorize Dr. Tom J. Pousti and such assistant as may be selected to perform the following procedure or treatment:

LIPOSUCTION SURGERY

I have received the following information sheet:

INFORMED CONSENT FOR LIPOSUCTION SURGERY

I recognize that during the course of the operation and medical treatment or anesthesia, unforeseen conditions may necessitate different procedures than those above. I therefore authorize the above physician and assistants of designees to perform such other procedures that are in the exercise of his or her professional judgment necessary and desirable. The authority granted under this paragraph shall include all conditions that require treatment and are not known to my physician at the time the procedure has begun.

I consent to the administration of such anesthetics considered necessary or advisable. I understand that all forms of anesthesia involve risk and possibility of complications, injury, and sometimes death.

I acknowledge that no guarantee has been given by anyone as to the results that may be obtained.

I consent to the photographing or televising of the operation or procedure to be performed, including appropriate portions of my body, for medical, scientific or educational purposes, provided my identity is not revealed by the pictures.

For purposes of advancing medical education, I consent to the admittance of observers to the operating room.

I consent to the disposal of any tissue, medical devices or body parts, which may be removed.

I authorize the release of my social security number to appropriate agencies for legal reporting and medical device registration if applicable.

IT HAS BEEN EXPLAINED TO ME IN A WAY THAT I UNDERSTAND:

- a) THE ABOVE TREATMENT OR PROCEDURE TO BE UNDERTAKEN
- b) THERE MAY BE ALTERNATIVE PROCEDURE OR METHODS OF TREATMENT
- c) THERE ARE RISKS TO THE PROCEDURE OR TREATMENT PROPOSED

Sign A or B Below

A. I consent to the treatment or procedure and the above listed items. I have been asked if I want more detailed explanation, but I am satisfied with the explanation and do not want more information.

Patient or person authorized to sign for patient

Date

B. I consent to the treatment or procedure and above listed items. I requested and received, in substantial detail, further explanation of the procedure or treatment, other alternative procedures or methods or treatment and information about the material risks or the procedure of treatment.

Patient or person authorized to sign for patient

Date